



# Bacitracin for Injection, USP

For IM Use

Rx only

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Bacitracin and other antibacterial drugs, Bacitracin should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.


## WARNING

Nephrotoxicity: Bacitracin in parenteral (intramuscular) therapy may cause renal failure due to tubular and glomerular necrosis. Its use should be restricted to infants with staphylococcal pneumonia and empyema when due to organisms shown to be susceptible to bacitracin. It should be used only where adequate laboratory facilities are available and when constant supervision of the patient is possible. Renal function should be carefully determined prior to and daily during therapy. The recommended daily dose should not be exceeded and fluid intake and urinary output maintained at proper levels to avoid kidney toxicity. If renal toxicity occurs the drug should be discontinued. The concurrent use of other nephrotoxic drugs, particularly streptomycin, kanamycin, polymyxin B, polymyxin E (colistin), and neomycin should be avoided.

## DESCRIPTION

Sterile Bacitracin USP is an antibiotic for intramuscular administration. Bacitracin is derived from cultures of *Bacillus subtilis* (Tracey). It is a white to pale buff, hygroscopic powder, odorless or having a slight odor. It is freely soluble in water; insoluble in acetone, chloroform, and ether. While soluble in alcohol, methanol, and glacial acetic acid, there is some insoluble residue. It is precipitated from its solutions and inactivated by many of the heavy metals.

The structural formula is:



The molecular formula is: C<sub>66</sub>H<sub>103</sub>N<sub>17</sub>O<sub>16</sub>S. Bacitracin is comprised of a polypeptide complex and Bacitracin A is the major component in this complex.

The molecular weight of Bacitracin A is 1422.71.

## CLINICAL PHARMACOLOGY

Bacitracin exerts pronounced antibacterial action *in vitro* against a variety of gram-positive and a few gram-negative organisms. However, among systemic diseases, only staphylococcal infections qualify for consideration of bacitracin therapy. Bacitracin is assayed against a standard and its activity is expressed in units, 1 mg having a potency of not less than 50 units.

Susceptibility plate testing: If the Kirby-Bauer method of disk susceptibility is used, a 10 unit bacitracin disk should give a zone of over 13 mm when tested against a bacitracin-susceptible strain of *Staphylococcus aureus*. Absorption of bacitracin following intramuscular injection is rapid and complete. A dose of 200 or 300 units/kg every 6 hours gives serum levels of 0.2 to 2 mcg/mL in individuals with normal renal function. The drug is excreted slowly by glomerular filtration. It is widely distributed in all body organs and is demonstrable in ascitic and pleural fluids after intramuscular injection.

## INDICATIONS AND USAGE

In accord with the statements in the "Warning Box" the use of intramuscular bacitracin is limited to the treatment of infants with pneumonia and empyema caused by staphylococci shown to be susceptible to the drug.

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Bacitracin and other antibacterial drugs, Bacitracin should be

used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

## CONTRAINDICATIONS

This drug is contraindicated in those individuals with a history of previous hypersensitivity or toxic reaction to it.

## WARNINGS

*Clostridium difficile* associated diarrhea (CDAD) has been reported with use of nearly all antibacterial agents, including Bacitracin for Injection and may range in severity from mild diarrhea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon leading to overgrowth of *C. difficile*.

*C. difficile* produces toxins A and B which contribute to the development of CDAD. Hypertoxin producing strains of *C. difficile* cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhea following antibiotic use. Careful medical history is necessary since CDAD has been reported to occur over two months after the administration of antibacterial agents.

If CDAD is suspected or confirmed, ongoing antibiotic use not directed against *C. difficile* may need to be discontinued. Appropriate fluid and electrolyte management, protein supplementation, antibiotic treatment of *C. difficile*, and surgical evaluation should be instituted as clinically indicated.

#### PRECAUTIONS

See "Warning Box" for precautions in regard to kidney toxicity associated with intramuscular use of bacitracin. Adequate fluid intake should be maintained orally, or if necessary, by parenteral method.

As with other antibiotics, use of this drug may result in overgrowth of nonsusceptible organisms, including fungi. If superinfection occurs, appropriate therapy should be instituted.

Prescribing Bacitracin in the absence of a proven or strongly suspected bacterial infection or a prophylactic indication is unlikely to provide benefit to the patient and increases the risk of the development of drug-resistant bacteria.

#### Information for Patients

Patients should be counseled that antibacterial drugs including Bacitracin should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When Bacitracin is prescribed to treat a bacterial infection, patients should be told that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed.

Skipping doses or not completing the full course of therapy may (1) decrease the effectiveness of the immediate treatment and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by Bacitracin or other antibacterial drugs in the future.

Diarrhea is a common problem caused by antibiotics which usually ends when the antibiotic is discontinued. Sometimes after starting treatment with antibiotics, patients can develop watery and bloody stools (with or without stomach cramps and fever) even as late as two or more months after having taken the last dose of the antibiotic. If this occurs, patients should contact their physician as soon as possible.

#### ADVERSE REACTIONS

**Nephrotoxic reactions.** Albuminuria, cylindruria, azotemia. Rising blood levels without any increase in dosage.

**Other reactions.** Nausea and vomiting. Pain at site of injection. Skin rashes.

#### DOSAGE AND ADMINISTRATION

TO BE ADMINISTERED INTRAMUSCULARLY ONLY

Infant dose: For infants under 2500 grams-900 units/kg/24 hours in 2 or 3 divided doses. For infants over 2500 grams-1,000 units/kg/24 hours, in 2 or 3 divided doses. Intramuscular injections of the solution should be given in the

upper outer quadrant of the buttocks, alternating right and left and avoiding multiple injections in the same region because of the transient pain following injection.

**Preparation of Solutions**—Should be dissolved in sodium chloride injection containing 2 percent procaine hydrochloride. The concentration of the antibiotic in the solution should not be less than 5,000 units per mL nor more than 10,000 units per mL.

Diluents containing parabens should not be used to reconstitute bacitracin; cloudy solutions and precipitate formation have occurred.

Reconstitution of the 50,000 unit vial with 9.8 mL of diluent will result in a concentration of 5,000 units per mL.

Solutions are stable for one week when stored in a refrigerator 2° to 8°C (36° to 46°F).

#### HOW SUPPLIED

Bacitracin for Injection, USP is supplied as follows:

NDC	Bacitracin for Injection, USP	Package Factor
25021-116-30	50,000 units per vial	1 vial per carton
25021-116-31	50,000 units per vial	10 vials per carton

#### Storage Conditions

Store the unreconstituted product in a refrigerator at 2° to 8°C (36° to 46°F).

#### LATEX-FREE

**Sterile, Nonpyrogenic, Preservative-free.**



Mfd. for SAGENT Pharmaceuticals  
Schaumburg, IL 60195 (USA)  
Made in India

©2009 Sagent Pharmaceuticals, Inc.

September 2009

1009302